5. REFERENCES

1. Ahmed, I. (2001). Hypotriglyceridemic and hypocholesterolemic effects of anti-diabetic *Momordica charantia* (Karela) fruit extract in streptozotocin-induced diabetic rats. Diabetes Research and Clinical Practice, 51(3), 155–161.

2. Ahmed, I., Adeghate, E., Sharma, A.K., Pallot, D.J., Singh, J. (1998). Effects of *Momordica charantia* fruit juice on islet morphology in pancreas of streptozotocin-diabetic rat. Diabetes Research and Clinical Practice, 40, 145-151.

3. Ali, L., Khan, A.K., Mamun, M.I., Mosihuzzaman, M., Nahar, N., Nur-e-Alam, M., Rokeya, B. (1993). Studies on hypoglycemic effects of fruit pulp, seed, and whole plant of *Momordica charantia* on normal and diabetic model rats, Planta Medicine, 59, 408–412.

4. Basch, E., Gabardi, S., Ulbricht, C. (2003). Bitter melon (*Momordica charantia*): a review of efficacy and safety. American Journal of Health and Systemic Pharmacology, 65, 356–359.

5. Bhise, S.B., Salunkhe, V.R., Wachasundar, N.P. (2009). Clinical Evaluation of Antidiabetic Polyherbal Ayurvedic Formulation- M*adhujeevan Churna* using S*tevia rebaudiana,* Indian Jounal of Pharmacy Practice, 2(2), 58-65.

6. Chen, J., Tian, R., Qiu, M., Lu, L., Zheng, Y., Zhang, Z. (2008). Trinorcucurbitane and cucurbitane triterpenoids from the roots of Momordica charantia, Phytochemistry, 69**,** 1043-1048.

7. Chen, Q., Li, E.T. (2005). Reduced adiposity in bitter melon (*Momordica charantia*) fed rats is associated with lower tissue triglyceride and higher plasma catecholamines, British Journal of Nutrition, 93, 747–754.

8. Choi, J., Lee, K.T., Jung, H., Park, H.S., Park, H.J. (2002). Anti-rheumatoid arthritis effect of the *Kochia scoparia* fruits and activity comparison of momordin Ic, its prosapogenin and sapogenin, Archives of Pharmacological Research, 25, 336–342.

9. Day, C., Cartwright, T., Provost, J., Bailey, C.J. (1990). Hypoglycaemic effect of *Momordica charantia* extracts, Planta Medica, 56, 426–429.

10. Dengiz, G.O., Gursan, N. (2005). Effects of *Momordica charantia* L. (Cucurbitaceae) on indomethacin-induced ulcer model in rats, Turkish Journal of Gastroenterology, 16, 85–88.

11. Giron, L.M., Freire, V., Alonzo, A. Caceres, A. (1991). Ethnobotanical survey of the medicinal flora used by the Caribs of Guatemala, Journal of Ethnopharmacology, 34, 173–187.

12. Grover, J.K., Yadav, S.P. (2004). Pharmacological actions and potential uses of *Momordica charantia*: a review, Journal of Ethnopharmacology, 93, 123-132.

13. Grover, J.K., Vats, V., Rathi, S.S., Dawar, R. (2001). Traditional Indian antidiabetic plants attenuate progression of renal damage in streptozotocin induced diabetic mice, Journal of Ethnopharmacology, 76, 233-238.

14. Gurbuz, I., Akyuz, C., Yesilada, E., Sener, B. (2000). Anti-ulcerogenic effect of *Momordica charantia* L. fruits on various ulcer models in rats, Journal of Ethnopharmacology, 71, 77–82.

15. Hayashi, K., Takehisa, T., Hamato, N., Takano, R., Hara, S., Miyata, T., Kato, H. (1994). Inhibition of serine proteases of the blood coagulation system by squash family protease inhibitors, Journal of Biochemistry, 116, 1013–1018.

16. Higashino, H., Suzuki, A., Tanaka, Y., Pootakham, K. (1992). Hypoglycemic effects of Siamese *Momordica charantia* and *Phyllanthus urinaria* extracts in

streptozotocin-induced diabetic rats (the 1st report), Nippon Yakurigaku Zasshi, 100, 415–421.

17. Husain, J., Tickle, I.J., Wood, S.P. (1994). Crystal structure of momordin, a type I ribosome inactivating protein from the seeds of *Momordica charantia*, FEBS Letter, 342, 154–158.

18. Jayasooriya, A.P. (2000). Effects of *Momordica charantia* powder on serum glucose levels and various lipid parameters in rats fed with cholesterol-free and cholesterol-enriched diets, Journal of Ethnopharmacology, 72 (1–2), 331–336.

19. Khan, M.R. (1998). *Momordica charantia* and *Allium sativum*: broad-spectrum antibacterial activity, Korean Journal of Pharmacology, 29,155-158.

20. Kohler, I., Jenett-Siems, K., Siems, K., Hernandez, M.A., Ibarra, R.A., Berendsohn, W.G., Bienzle, U. Eich, E. (2002). In vitro antiplasmodial investigation of medicinal plants from El Salvador, Zeitschrift fur Naturforschung [section-C], 57, 277–281.

21. Lans, C., Brown, G. (1998). Observations on ethno veterinary medicines in Trinidad and Tobago, Preventive Veterinary Medicine, 35, 125–142.

22. Lee-Huang, S. (1995). Inhibition of the integrase of human immunodeficiency virus (HIV) type 1 by anti-HIV plant proteins MAP30 and GAP31, Proc Natl Acad Sci**,** 92(19), 818–22.

23. Limtrakul, P., Khantamat, O., Pintha, K. (2004). Inhibition of P-glycoprotein activity and reversal of cancer multidrug resistance by *Momordica charantia* extract, Cancer Chemotherapy Pharmacology, 54, 525–530.

24. Lolitkar, M.M., Rajarama, Rao, M.R. (1966). Pharmacology of a Hypoglycaemic Principles Isolated from the Fruits of *Momordica charantia* Linn, 28(5), 129-133.

25. Mastuda, H.L., Murakami, T., Matsumura, N., Yamahara, J., Yoshikawa, M. (1998). Antidiabetic principles of natural medicines, Chemical and Pharmaceutical Bulletin, 46, 1399-1403.

26. Matsuda, H., Li, Y., Yoshikawa, M. (1999). Roles of capsaicin-sensitive sensory nerves, endogenous nitric oxide, sulfhydryls, and prostaglandins in gastroprotection by momordin Ic, an oleanolic acid oligoglycoside, on ethanol-induced gastric mucosal lesions in rats, Life Science, 65, 27–32.

27. Miura, T., Itoh, C., Iwamoto, N., Kato, M., Kawai, M., Park, S.R., Suzuki, I. (2001). Hypoglycemic activity of the fruit of the *Momordica charantia* in type 2 diabetic mice, Journal of Nutrition Sciences Vitaminology, 47, 340–344.

28. Modak. M., Dixit, P., Londhe, J., Ghaskadbi, H., Devasagayam, T. (2007). Indian Herbs and Herbal Drugs used for treatment of Diabeties, J Clin Biochem Nutr, 40, 163–173.

29. Murakami, T., Emoto, A., Matsuda, H., Yoshikawa, M. (2001). Medicinal Foodstuffs, XXI, Structures of New Cucurbitaceae-Type Triterpene Glycosides, Chem Pharm Bull, 49, 54-63.

30. Nakamura, S., Murakami, T., Nakamura, J., Kobayashi, H., Matsuda, H. Yoshikawa, M. (2006). Structures of New Cucurbitane Type Triterpenes and Glycosides, Karavilagenins and Karavilosides, from dried fruit of *Momordica charantia* Linn. In Sri Lanka, Chem Pharm Bull, 54, 1545-1550.

31. Ng, T.B., Chan, W.Y., Yeung, H.W. (1992). Proteins with abortifacient, ribosome inactivating, immunomodulatory, antitumor and anti-AIDS activities from Cucurbitaceae plants, General Pharmacology, 23, 579– 590.

32. Ojewole, J.A., Adewole, S.O., Olayiwola, G. (2006). Hypoglycaemic and hypotensive effects of Momordica charantia Linn (Cucurbitaceae) whole-plant aqueous extract in rats, Cardiovascular Journal of South Africa, 17**,** 227-232.

33. Omoregbe, R.E., Ikuebe, O.M., Ihimire, I.G. (1996). Antimicrobial activity of some medicinal plants extracts on *Escherichia coli*, *Salmonella paratyphi* and *Shigella dysenteriae*, African Journal of Medical Science, 25, 373–375.

34. Parkash, A., Ng, T.B., Tso, W.W. (2002) Purification and characterization of charantin, a napin-like ribosome-inactivating peptide from bitter gourd (*Momordica charantia*) seeds, Journal of Peptide Research, 59, 197–202.

35. Patel, P.M., Patel, K.N., Patel, N.M., Goyal, R.K. (2006). Development of HPTLC method for estimation of charantin in herbal formulations, Pharmacognosy Magazine, 2(8), 224-226.

36. Patel, S.S., Shah, R.S., Goyal, R.K. (2009). Antihyperglycemic, Antihyperlipidemic and antioxidant effects of Dihar, a polyherbal Ayurvedic formulation in streptozotocin induced diabetic rats, Indian Journal of Experimental Biology, 47, 564-570.

37. Raman, A., Lau, C. (1996). Anti-diabetic properties and phytochemistry of *Momordica charantia* L. (Cucurbitaceae), Phytomedicine, 2, 349–362.

38. Ross, I.A. (1999). Medicinal Plants of the World, Humana Press, New Jersey, USA, 213–219.

39. Sathishsekar, D., Subramanian, S. (2005a). Beneficial effects of *Momordica charantia* seeds in the treatment of STZ-induced diabetes in experimental rats, Biological and Pharmaceutical Bulletin, 28, 978–983.

40. Sathishsekar, D., Subramanian, S. (2005b). Antioxidant properties of *Momordica Charantia* (bitter gourd) seeds on streptozotocin induced diabetic rats. Asia Pacific Journal of Clinical Nutrition, 14, 153–158.

41. Schmourlo, G., Mendonca-Filho, R.R., Alviano, C.S., Costa, S.S. (2005). Screening of antifungal agents using ethanol precipitation and bioautography of medicinal and food plants, Journal of Ethnopharmacology, 96, 563–568.

42. Senanayake, V.K., Maruyama, M., Shibuya, K., Masanobu, S., Fukuda, N., Morishita, T. (2004). The effects of bitter melon (*Momordica charantia*) on serum and liver triglyceride levels in rats, Journal of Ethnopharmacology, 91, 257-262.

43. Shibib, B.A., Khan, L.A., Rahman, R. (1993). Activity of *Coccinia indica* and *Momordica charantia* in diabetic rats: depression of the hepatic gluconeogenic enzymes glucose-6-phosphatase and fructose-1,6- bisphosphatase and elevation of both liver and red-cell shunt enzyme glucose-6-phosphate dehydrogenase, Biochemistry Journal, 292, 267–270.

44. Shih, C.C., Lin, C.H., Lin, W.L. (2008). Effects of Momordica charantia on insulin resistance and visceral obesity in mice on high-fat diet, Diabetes Research and Clinical Practice, 81**,** 134-43.

45. Sun, Y., Huang, P.L., Li, J.J., Huang, Y.Q., Zhang, L., Huang, P.L., Lee-Huang, S. (2001). Anti-HIV agent MAP30 modulates the expression profile of viral and cellular genes for proliferation and apoptosis in AIDS-related lymphoma cells infected with Kaposi's sarcomaassociated virus, Biochemical and Biophysical Research Communication, 287, 983–994.

46. Terenzi, A., Bolognesi, A., Pasqualucci, L., Flenghi, L., Pileri, S., Stein, H., Kadin, M. (1996). Anti-CD30 (BER=H2) immunotoxins containing the type-1 ribosome-inactivating proteins momordin and PAP-S (pokeweed antiviral protein from seeds) display powerful antitumour activity against CD30+ tumour cells in vitro and in SCID mice, British Journal of Haematology, 92, 872–879.

47. Virdi, J., Sivakami, S., Shahani, S., Suthar, A.C., Banavalikar, M.M., Biyani, M.K. (2003). Antihyperglycemic effects of three extracts from Momordica charantia. Journal of Ethnopharmacology, 88**,** 107-111.

48. Yesilada, E., Gurbuz, I., Shibata, H. (1999). Screening of Turkish antiulcerogenic folk remedies for anti-*Helicobacter pylori* activity, Journal of Ethnopharmacology, 66, 289–293.

49. Yuan, Y.R., He, Y.N., Xiong, J.P., Xia, Z.X. (1999). Three-dimensional structure of beta-momorcharin at 2.55 A resolution, Acta Crystallographica Section D-Biological Crystallography, 55, 1144–1151.

50. Zhen, Z.X., Teng, J.Y., Liu, J.Y., Qiu, J.H., Ouyang, H., Xue, C. (2005). The hypoglycemic effects of crude polysaccharides extract from *Momordica charantia* in mice, Wei Sheng Yan Jiu, 34, 361–363.